

1

The Dual Quark Models ^a

K. Bardakci and M.B.Halpern

*Department of Physics, University of California,
Theoretical Physics Group, Lawrence Berkeley National Laboratory
University of California, Berkeley CA 94720 USA*^a To appear in the volume “The Birth of String Theory”, Ed. P. Di Vecchia (Cambridge University Press, Cambridge UK)**Abstract**

We briefly recall the historical environment around our 1971 and 1975 constructions of current-algebraic internal symmetry on the open string. These constructions included the introduction of world-sheet fermions, the independent discovery of affine Lie algebra in physics (level one of affine $su(3)$), the first examples of the affine-Sugawara and coset constructions, and finally – from compactified spatial dimensions on the string – the first vertex-operator constructions of the fermions and level one of affine $su(n)$.

1.1 Introduction

In this contribution, we describe the environment around our 1971 paper “New Dual Quark Models” [BH71] and the two companion papers “The Two Faces of a Dual Pion-Quark Model” [Hal71c] and “Quantum Solitons which are $SU(N)$ Fermions” [Hal75] in 1971 and 1975, which laid the foundations of non-Abelian current-algebraic internal symmetry on the string.

The background for our contributions included many helpful discussions with other early workers, including H.M.Chan, C. Lovelace, H. Ruegg and C. Schmidt (during KB’s 1969 visit to CERN) as well as R. Brower, S. Klein, C. Thorn, M. Virasoro and J. Weis (with MBH at Berkeley). Both of us also acknowledge many discussions with Y. Frishman, G. Segre, J. Shapiro and especially S. Mandelstam. Later discussions with I. Bars and W. Siegel are also acknowledged by MBH. With apologies to many other authors then, we reference here only the work which was most influential in our early thinking: before Veneziano, there had been widespread interest in the quark model of Gell-Mann and Zweig, including the four-dimensional current algebra (Adler and Dashen [AD68]) of quarks.

1.2 The Bosonic Dual Model

In 1970, the work of Fubini and Veneziano [FV69] together with the Virasoro algebra [Vir70] provided an *algebraic formulation* of open bosonic string theory which we found quite exciting. The first ingredients of this formulation are the bosonic oscillators

$$\begin{aligned} [J_\mu(m), J_\nu(n)] &= -m g_{\mu\nu} \delta_{m+n,0}, \\ m, n \in \mathbb{Z}, \quad g &= \text{diag}(1, -1, \dots, -1), \quad \mu, \nu = 0, 1, \dots, 25 \end{aligned} \quad (1.1)$$

which we have rescaled here as the modes of an Abelian current algebra on the circle. The conformal-field-theoretic convention for the local currents associated to these modes are

$$J_\mu(z) = \sum_{m \in \mathbb{Z}} J_\mu(m) z^{-m-1} = i \partial_z Q_\mu(z) \quad (1.2)$$

where

$$Q(z) = q - iJ(0)\ln(z) + i \sum_{m \neq 0} J(m) \frac{z^{-m}}{m} \quad (1.3)$$

are the Fubini-Veneziano fields and $\{q\}$ is canonically conjugate to the momenta $\{p = J(0)\}$. Normal-ordered exponentials of the Fubini-Veneziano fields are the string vertex operators, to which we shall return below.

In our discussions we shall emphasize the mode algebras, but for various modern applications the complex variable z in the local fields can be taken in many forms, including $\exp(i\theta)$ (the circle or loop), $\exp(i(\tau + \sigma))$ (left-movers on a closed string), the Euclidean version $\exp(\tau + i\sigma)$ for conformal field theory (CFT), or the real line for open strings. These alternate forms led to some confusion in the early days of string theory, but in fact most of the early work reviewed in this article was in the context of the open string, using the parametrization $z = \exp(i\theta)$ on the circle.

Finally, the Virasoro algebra of the critical open bosonic string is

$$\begin{aligned} L(m) &= -\frac{1}{2} g^{\mu\nu} \sum_{n \in \mathbb{Z}} : J_\mu(n) J_\nu(m-n) : \\ [L(m), L(n)] &= (m-n) L(m+n) + \frac{26}{12} m(m^2-1) \delta_{m+n,0} \end{aligned} \quad (1.4)$$

where the symbol $:\dots:$ is bosonic normal-ordering. Under these generators the Abelian currents $\{J(z)\}$ covariantly transform with conformal weight equal to one.

Taken together, the algebraic statements in Eqs. (1.1)-(1.3) provided the first example of modern CFT (see Belavin, Polyakov and Zamolodchikov

[BPZ84]). We have here taken an historical liberty, because the central extension of the Virasoro algebra was not observed until 1974, and even then only in a private communication from J. Weis quoted in Ref. [CT74] (Chodos and Thorn). Throughout this article we adopt CFT conventions, including an extra factor $z^{-\Delta}$ (see Eq. (1.1)) for the mode expansions of Virasoro primary fields with conformal weight Δ . The CFT convention guarantees that a given Virasoro primary field creates a Virasoro primary state of the same conformal weight as z goes to 0 on the vacuum (in this case the zero-momentum vacuum $|\{p = J(0) = 0\}\rangle$ of the open bosonic string). On the other hand, the extra factor $z^{-\Delta}$ must be removed for exact correspondence of the local fields with $z = \exp(i\theta)$ in early work on the open string.

1.3 World-Sheet Fermions

As it appeared then, the open bosonic string had two deficiencies beyond the tachyonic ground state. In the first place Lie-algebraic internal symmetries were implemented only by the multiplicative Chan-Paton factors [PC69], which live on the boundary of the string. Second, the bosonic string had no spin, that is, no space-time fermions. Our first attempt to solve the second problem was a generalization of Chan-Paton factors to include spin [BH69], but this multiplicative approach was unsuccessful because it included negative-norm states.

It seemed natural then to approach the problems of spin and internal symmetry in a different way, by introducing new local “quark” fields on the string – now called *world-sheet fermions* [BH71]:

$$\bar{\psi}^I(z) = \sum_{p \in \mathbb{Z} + \frac{1}{2}} \bar{\psi}^I(p) z^{-p - \frac{1}{2}}, \quad \psi_I(z) = \sum_{p \in \mathbb{Z} + \frac{1}{2}} \psi_I(p) z^{-p - \frac{1}{2}} \quad (1.5)$$

whose modes obey anticommutation relations

$$\begin{aligned} [\psi_I(q), \psi_J(p)]_+ &= [\bar{\psi}^I(q), \bar{\psi}^J(p)]_+ = 0, \\ [\psi_I(q), \bar{\psi}^J(p)]_+ &= \delta_I^J \delta_{q+p,0}, \quad q, p \in \mathbb{Z} + \frac{1}{2}. \end{aligned} \quad (1.6)$$

As in the four-dimensional quark model itself, the generic indices $\{I, J\}$ were designed so that the world-sheet fermions could be space-time spinors $\{\psi_r\}$ or carry internal symmetry or both $\{\bar{\psi} \gamma_\mu \lambda_a \psi\}$. The Virasoro generators of our world-sheet fermions are

$$L(m) = \sum_{p \in \mathbb{Z} + \frac{1}{2}} \left(p + \frac{m}{2} \right) : \bar{\psi}^I(p) \psi_I(m - p) : \quad (1.7)$$

with fermionic normal-ordering and central charge c equal to the number of $(\psi, \bar{\psi})$ pairs. In what follows, we often refer to our world-sheet fermions as the *dual quarks*.

Under these Virasoro generators, the dual quark fields $\{\psi, \bar{\psi}\}$ are Virasoro primary fields with conformal weight $1/2$ and trivial monodromy. The half-integer moding of the dual quark fields guaranteed a unique vacuum (or ground state) $|0\rangle$ which is annihilated by the positive modes of both fields, and the adjoint operation is defined for internal symmetry as $\psi(p)^\dagger = \bar{\psi}(-p)$ (with a γ_0 modification for spin). In either case, our Virasoro generators satisfy the generalized hermiticity $L(m)^\dagger = L(-m)$ and the vacuum state conditions:

$$L(m)|0\rangle = 0, \quad m \geq -1. \quad (1.8)$$

With internal symmetry only, our dual quark models then provided the first examples of *unitary* CFT, with a positive-definite Hilbert space on a non-degenerate vacuum with conformal weight $\Delta = 0$. In modern parlance, the half-integer moded complex BH fermions are related to anti-periodic Weyl fermions on the world sheet, but it should be born in mind that fermionic world-sheet action formulations of open string theory had not been studied at that time.

Were these the first world-sheet fermions? In fact, our work was essentially simultaneous with that of Ramond [Ram71]: Our paper was submitted in November of 1970 while Ramond's paper with integer-moded Majorana-Weyl world-sheet fermions was submitted in January 1971; the two papers were published in the same issue of Physical Review. As we now know the R fermions $\{b_\mu\}$, with their somewhat-counterintuitive space-time vector label μ and superconformal symmetry, provided the correct covariant description of space-time fermions and – in a perfect match to the R construction – half-integer moded Majorana-Weyl world-sheet fermions $\{b_\mu\}$ were later introduced for the bosonic sector by Neveu and Schwarz [NS71].

For the historical record, the lure of our seemingly more physical space-time spinor fields $\{\psi_r\}$ persisted for some time, leading to many connections (Halpern [Hal71a], [Hal71b], [Hal71c], Bardakci [Bar71], Halpern and Thorn [HT71a], [HT71b] and Mandelstam [Man73a], [Man73b]) between our fields and those of the RNS formulation (We mention in particular the first example of half-integer-moded scalar fields in [HT71b]). Our original intuition was however not realized until 1982, when a world-sheet fermion with 10-dimensional Majorana-Weyl spinor indices was successfully incorporated by Green and Schwarz in the light-cone formulation of superstring theory [GS82].

In what follows, we therefore restrict the discussion primarily to our various current-algebraic and CFT constructions on the open string.

1.4 Affine Lie algebra

For continuity with the bosonic string, we begin this discussion with a footnote of our 1971 paper – which observed that the spatial components $\{I = 1, \dots, 25\}$ of the Abelian currents (1.1),(1.2) could be constructed out of our world-sheet fermions [BH71]:

$$\begin{aligned} J_I(m) &= \sum_{p \in \mathbb{Z} + \frac{1}{2}} : \bar{\psi}^I(p) \psi_I(m-p) :, \\ J_I(z) &= : \bar{\psi}^I(z) \psi_I(z) : = i \partial_z Q_I(z). \end{aligned} \quad (1.9)$$

This result is now recognized as a simple but important part of Bose-Fermi equivalence on the open string, a subject to which we will return below.

Our next construction was the independent discovery of affine Lie algebra in physics [BH71], still sometimes called the “Dual Quark Model”, in which the non-abelian currents are naturally constructed as dual-quark bilinears with conformal weight 1:

$$J_a(z) = \sum_{m \in \mathbb{Z}} J_a(m) z^{-m-1} = : \bar{\psi}(z) T_a \psi(z) : \quad (1.10)$$

$$\begin{aligned} J_a(m) &= \sum_{p \in \mathbb{Z} + \frac{1}{2}} : \bar{\psi}^I(p) (T_a)_I^J \psi_J(m-p) :, \\ [T_a, T_b] &= i f_{ab}^c T_c, \quad \text{Tr}(T_a T_b) = k \eta_{ab}. \end{aligned} \quad (1.11)$$

Here $\{T\}$ can be any irreducible matrix representation of any simple Lie algebra g , with structure constants $\{f\}$ and Killing metric $\{\eta\}$. Then one finds that the modes of the dual quark-model currents $\{ : \bar{\psi} T \psi : \}$ are the generators of the *affine Lie algebra*

$$\begin{aligned} [J_a(m), J_b(n)] &= i f_{ab}^c J_c(m+n) + m (k \eta_{ab}) \delta_{m+n,0}, \\ a, b, c &= 1 \dots \dim(g), \quad m, n \in \mathbb{Z} \end{aligned} \quad (1.12)$$

where the quantity k in the central term is called the *level* of the affine algebra. (We assume in this article that the highest root-length squared of Lie g is two, in which case the level in the dual quark models is also equal to the Dynkin index of irreducible representation T .) Affine Lie algebra is also known as centrally-extended non-Abelian current algebra on the circle. In this construction, the dual-quark fields $\{\psi, \bar{\psi}\}$ transform respectively in

the $\{T, \bar{T}\}$ representations of Lie g – the Lie algebra being generated by the zero modes $\{J_a(0)\}$ of the affine algebra.

In fact our first example of this construction used the 3×3 Cartesian Gell-Mann matrices $\{T_a \propto \lambda_a\}$ of $su(3)$ (and the Dynkin index of the fundamental of $su(n)$ is 1) – so the *first concrete representation of affine Lie algebra* was $su(3)_1$, that is, level one of affine $su(3)$ (see also Sec. 1.8). In this case, as in the four-dimensional quark model itself, the dual-quark fields $\{\psi, \bar{\psi}\}$ transform respectively in the $(3, \bar{3})$ representations of $su(3)$. Dual quark-model constructions of the currents of $so(3, 1)_1$ and $so(4, 1)_1$ were also included in our 1971 paper, and further studied along with those of higher Lorentz algebras in the first companion paper [Hal71c]. The dual quark models of $su(n)_1$ were given later in the second companion paper [Hal75].

Because connections between open strings and field theory were not understood at the time, our interpretation of the central terms in our string-theoretic constructions of affine Lie algebras on the circle was based on the presence of such a term in the Abelian mode algebra (1.1) of the open bosonic string – to which our constructions reduced in the Abelian case. This understanding was supported by our fermionic construction (see Eq. (1.7) above) of the modes of the Abelian current algebra.

On the other hand, we often think now of the central term in affine Lie algebra as an analogue on the circle of the equal-time Schwinger term [Sch59] (indeed, see Eq. (3.10) in Ref. [BH71]), so we briefly sketch the history of $su(n)$ current algebra in the two-dimensional Lorentz-invariant quantum field theory of massless free fermions: The equal-time Schwinger term in $su(n)$ current algebra on the line was first computed in 1969 (see Coleman, Gross and Jackiw [CGJ69]), and the one-dimensional $su(n)$ current algebras of the light-cone currents $\{J_a^\pm(t \pm x)\}$ – with one-dimensional Schwinger terms – were given later by Dashen and Frishman [DF73]. In fact however the momenta are continuous in this non-compactified, Lorentz-invariant context, so our open-string mode forms of the fermions and affine Lie algebra on the circle [BH71] do not appear in these papers.

For a modern introduction to affine Lie algebras in CFT, see the work by Gepner and Witten [GW86] and also the representation theory for all integer levels in Kac [Kac90]. In physics the affine algebras have sometimes also been called Kac-Moody algebras, but this term is more properly reserved for the general algebraic systems including affine and hyperbolic algebras (see Sec. 1.8).

1.5 The Affine-Sugawara Constructions

The *affine-Sugawara constructions* [BH71] [Hal71c] are the simplest realization of Virasoro generators as quadratic forms on the generators of affine Lie algebras ,

$$\begin{aligned} L_g(m) &= \frac{\eta^{ab}}{2k + Q_g} \sum_{n \in \mathbb{Z}} : J_a(n) J_b(m - n) :, \\ [L_g(m), L_g(n)] &= (m - n) L_g(m + n) + \frac{c_g}{12} m(m^2 - 1) \delta_{m+n,0}, \\ c_g &= \frac{2k \dim(g)}{2k + Q_g}. \end{aligned} \quad (1.13)$$

The quantity Q_g is the quadratic Casimir of Lie g , which in our present root-normalization is twice the dual Coxeter number of g . The original Sugawara model [Sug68], Sommerfield [Som68], Bardakci, Frishman and Halpern [BFH68], Bardakci and Halpern [BH68], Gross and Halpern [GH69], Coleman, Gross and Jackiw [CGJ69] was in four dimensions on a different current algebra, the so-called algebra of fields. The affine-Sugawara (or Sugawara-like) constructions in Eq. (1.11) hold for all integer levels of any affine Lie algebra.

The first example of the affine-Sugawara construction [BH71] used the $su(3)_1$ dual quark-model currents of the previous section (the dual Coxeter number of $su(n)$ is n). The affine-Sugawara construction on the level-one currents of affine $so(4,1)$ was also given in our 1971 paper, and extended in the first companion paper [Hal71c] to the construction on the level-one currents of affine $so(3,1)$ and higher Lorentz algebra. The case of $su(n)_1$ was given later in [Hal75]. We emphasize that the prefactors $(2k + Q_g)^{-1}$ in these constructions follow from careful consideration of the normal ordering $: \dots :$ of the current modes. The affine-Sugawara constructions, being quartic in our dual quarks, were called “current-current” or “spin-spin” interactions in those days, and models which used the dual quarks or the affine-Sugawara constructions instead of extra dimensions on the open string were known as “additive” models.

In physics the general form (1.11) of the affine-Sugawara construction was first given by Knizhnik and Zamolodchikov [KZ84] thirteen years after our examples, and used by these authors to find the general KZ equations. In the same year, the important paper by Witten [Wit84] gave the action formulation and non-Abelian bosonization of these operator systems, leading to their understanding – especially at higher level – as interacting quantum field theories of the WZW type [GW86]. The independent history of the affine-Sugawara constructions in mathematics is noted in Sec. 1.8.

In this connection, we also sketch the history of the $u(n)$ Sugawara-like constructions in two-dimensional Lorentz-invariant field theories of massless free fermions: Using a point-splitting regularization of the products of $u(n)$ currents, Ref. [CGJ69] showed the equivalence of the two-dimensional analogue $\{T_{\mu\nu}\}$ of the original Sugawara stress-tensor to the free-fermionic stress-tensor $\{\Theta_{\mu\nu}\}$, and (building on the Abelian case in Ref. Dell’Antonio, Frishman and Zwanziger [DFZ72]) normal-ordered light-cone analogues of the $u(n)$ Sugawara construction were later given in Ref. [DF73] (Eqs. (5.15)-(5.16) of Ref. [DF75] also gave (and solved) light-cone analogues of the $u(n)$ KZ equations for the fermionic four-point function.). Again however, these non-compactified, Lorentz-invariant results had continuous momenta, so the corresponding affine-Sugawara constructions do not appear in these papers. (See also Refs. [Hal75], [BHN76]).

1.6 The Coset Constructions

The affine-Sugawara construction (1.13) on the currents of affine $\mathfrak{g}$ is invariant under Lie $\mathfrak{g}$, so we moved next to the study of *symmetry breaking* in current-algebraic CFT on the string.

We begin this discussion with the results in our paper [BH71]. In particular for our example on $su(3)_1$, we first studied the addition to the affine-Sugawara construction of a term $\lambda J_8(m) + \lambda^2 \delta_{m,0}$ for all constant λ , which gave Virasoro generators with *continuous* $su(3)$ -symmetry breaking. This was the first example of what is today known as inner-automorphic twists or c-fixed *conformal deformations* (Freericks and Halpern [FH88]).

Next we looked for $su(3)$ -symmetry breaking solutions on $su(3)_1$ when the Virasoro generators were assumed to have the more general quadratic form $L = \sum_a C_a : J_a J_a :$ with constant coefficients $\{C_a, a = 1 \dots 8\}$. We stated that solutions beyond the affine-Sugawara construction for the coefficients $\{C_a\}$ existed, giving Virasoro generators with *quantized* $su(3)$ -symmetry breaking. This was the first *implicit* mention of coset constructions, but we gave no details about these solutions – noting only that they all had “infinite-degeneracy” problems. This issue (amplified in our non-additive “spin-orbit” constructions of the next section of this reference) was that the solutions to the Virasoro conditions come in *commuting* “K-conjugate pairs” of Virasoro generators $[L(m), \tilde{L}(n)] = 0$ which sum to the affine-Sugawara construction $L_g(m) = L(m) + \tilde{L}(m)$. This phenomenon was called *K-conjugation covariance* [BH71], [Hal71c], [Man73a], [Man73b], and the associated spectral degeneracy of each commuting K-conjugate partner, due to the other, was called K-degeneracy.

The first *explicit* examples of non-Abelian coset constructions were given in the companion paper “The Two Faces of a Dual Pion-Quark Model” [Hal71c], which solved the Virasoro conditions for all possible (BH quartic) conformal spin-spin interactions among the four-dimensional dual quark bilinears T,V,A,P and S ($\{T_{\mu\nu} =: \bar{\psi}\sigma_{\mu\nu}\psi : \}$ and so on). All 31 of these conformal constructions were affine-Sugawara or coset constructions, and a number of these were included explicitly in the applications of the text (see Eqs. (3.7)(3.9) and (3.14)). Each of these results was presented as a decomposition of an affine-Sugawara construction into a commuting K-conjugate pair of conformal constructions – which we would today write as

$$L_g(m) = L_{g/h}(m) + L_h(m) , \quad [L_{g/h}(m), L_h(n)] = 0 , \quad (1.14)$$

where g is the original symmetry algebra, h is any subalgebra of g and g/h is the coset space.

We will focus here on the Lie-algebraic identifications in the simplest case, given as $N^{NS} + N^5 = N^H$ in Eq. (3.14) of the paper. All three sets of Virasoro generators are defined in Eqs. (3.9) and (3.14) as different linear combinations of the BH quartics T^2 and A^2 , and it is not difficult to see that that N^{NS} and N^H are affine-Sugawara constructions respectively on $so(3,1)_1$ and $so(4,1)_1$. Therefore, Eq. (3.14) can be read as the K-conjugate decomposition

$$L_{so(3,1)}(m) + L_{so(4,1)/so(3,1)}(m) = L_{so(4,1)}(m) \quad (1.15)$$

into the commuting K-conjugate partners on $so(3,1)$ and $so(4,1)/so(3,1)$. In fact, this result was checked explicitly with the identities given in the text between the BH dual-quark bilinears T,A and certain five-dimensional NS bilinears (see Eq. (2.9) of the text). Using these identities, it was noted that each quartic BH Virasoro generator collapses to an NS bilinear, the affine-Sugawara constructions N^{NS} and N^H on $so(3,1)_1$ and $so(4,1)_1$ being equal respectively to the four- and five-dimensional quadratic NS Virasoro generators, while the non-abelian coset construction N^5 (with central charge $c_{so(4,1)/so(3,1)} = 5/2 - 2 = 1/2$) is equal to the quadratic Virasoro generators of the fifth NS operator alone. Similar quartic-to-bilinear equivalences had been seen earlier in [CGJ69], [BH71].

The general form of the coset construction for all $h \subset g$

$$L_{g/h}(m) = L_g(m) - L_h(m), \quad c_{g/h} = c_g - c_h \quad (1.16)$$

was given by Goddard, Kent and Olive [GKO85] fourteen years after our examples. The K-degeneracy of the Virasoro construction on g/h is now understood as the local gauge invariance associated to the h-currents of the

commuting K-conjugate affine-Sugawara construction $\{L_h(m)\}$ on $\mathfrak{h}$. It should be emphasized however that the coset constructions are by no means the end of CFT constructions on the currents of affine Lie algebras: Our original method of solving for the coefficients $\{L^{ab}\}$ in quadratic forms $\{L^{ab} : J_a J_b : \}$ eventually led to the completely K-conjugation-covariant *Virasoro master equation* (Halpern and Kiritsis [HK89]), which includes the affine-Sugawara constructions and the coset constructions as very special cases (with rational central charge) of irrational conformal field theory (Halpern, Kiritsis, Obers and Clubok [HKOC96]).

1.7 The Vertex-Operator Constructions

There is one more important strand which belongs in this story – namely the second companion paper in 1975 “Quantum Solitons which are SU(N) Fermions” [Hal75], which described –using compactified spatial dimensions on the open string – the first vertex-operator constructions of world-sheet fermions and affine Lie algebra.

By 1972, interest among physicists was shifting back to field theory. We mention in particular the Bose-Fermi equivalence of the sine-Gordon and interacting Thirring model studied by Coleman [Col75] and Mandelstam [Man75] in 1975. (See also the free-field bosonization of the single fermion bilinears by Kogut and Susskind [KS75].) One of us (MBH) realized that – with two observations – this development could be applied to our 1971 non-Abelian current-algebraic constructions on the string.

The first observation was that Mandelstam’s normal-ordered line-integral construction of a single interacting fermion could be easily generalized to *many* interacting fermions on the line by including appropriate Klein transformations [Kle38]. This led to bosonic realizations of centrally-extended $su(n)$ current algebras on the line, and the equivalence of various generalized sine-Gordon models with corresponding sets of interacting $u(n)$ Thirring models. In particular, the $su(n)$ currents were constructed from $(n-1)$ independent two-dimensional bosons, showing the roots of $su(n)$ in the off-diagonal currents. (See also the extension of these ideas to the bosonization of two-dimensional non-abelian gauge theories [Hal76] – and in particular Eq.(3.13) of this reference.)

The second observation in the 1975 paper was that, for massless free fermions and bosons, there was a structural parallel between Mandelstam’s fermions on the line and particular string vertex operators on the circle. Indeed, a formal projective map (from the line to the circle) was given between each two-dimensional left- and right- mover free-field construction

in the field theories and a *pair* of our open-string counterparts on the circle, a structure which corresponded in fact to closed strings. These points were emphasized in the second appendix “Connections with Dual Models” of that paper – which also gave the vertex-operator construction of many world-sheet fermions [Hal75]

$$\bar{\psi}^I(z) = : \exp(i Q_I(z)) : \bar{\xi}^I, \quad \psi_I(z) = \xi_I : \exp(-i Q_I(z)) :. \quad (1.17)$$

where $\{Q_I\}$ are the Fubini-Veneziano fields and $\{\bar{\xi}, \xi\}$ are the Klein transformations of the text. This relation guaranteed that the world-sheet fermions were Virasoro primary fields with conformal weight $1/2$ under the open-string bosonic Virasoro generators, and we identified these constructions as our half-integer-moded dual quarks [BH71]. Although it was not explicitly mentioned, this identification had in fact been checked from vacuum expectation values using the natural identification of the bosonic zero-momentum vacuum $|\{p = J(0) = 0\}\rangle$ as the unique vacuum $|0\rangle$ annihilated by the positive modes of the dual quarks.

The vertex-operator construction of $su(n)_1$ can therefore be obtained immediately in a single step by direct substitution of the fermionic vertex-operators (1.14) into our dual quark-model construction $\{J_a \propto : \bar{\psi} \lambda_a \psi : \}$ of the currents. We regret having omitted this last simple step in the appendix, thinking it implicit in the context of the paper: The off-diagonal currents $: \bar{\psi}^I(z) \psi_J(z) : \propto : \exp(i(Q_I(z) - Q_J(z))) : , I \neq J = 1 \dots n$ of $su(n)_1$ follow immediately, showing the roots of $su(n)$ on $(n-1)$ independent fields, and the diagonal currents of $su(n)_1$ are a derivative of the same $(n-1)$ fields, while the extra $u(1)$ current is a derivative of the normalized orthogonal sum $Q_+(z) = (1/\sqrt{n}) \sum_I Q_I(z)$. We state here the result only for $su(2)_1 + u(1)$

$$\begin{aligned} J_{\pm}(z) &= : \bar{\psi}(z) \tau_{\pm} \psi(z) : = : \exp(\pm i \sqrt{2} Q_{\pm}(z)) :, \\ J_3(z) &= \frac{1}{\sqrt{2}} : \bar{\psi}(z) \tau_3 \psi(z) : = i \partial_z Q_-(z), \\ J_0(z) &= \frac{1}{\sqrt{2}} : \bar{\psi}(z) \psi(z) : = i \partial_z Q_+(z), \\ Q_{\pm}(z) &= \frac{1}{\sqrt{2}} (Q_1(z) \pm Q_2(z)) \end{aligned} \quad (1.18)$$

which is in fact the only case where (as discussed in the text) the Klein transformations can be chosen to cancel. The charged currents $\{J_{\pm}(z)\}$ of $su(2)_1$ show the roots of $su(2)$, and results for the neutral current $J_3(z)$ and the extra $u(1)$ current $J_0(z)$ are of course closely related to our earlier Bose-Fermi relation on the string given in Eq. (1.7) above.

We concluded that we had constructed a current-algebraic $su(2)$ from a

single compactified spatial dimension, and also that the paper was equivalent to constructing as much as current-algebraic $u(22) = su(22) + u(1)$ from the compactified extra spatial dimensions of the critical open bosonic string. (Quantized extra momenta were well-known in early open-string theory). In the following year, Banks, Horn and Neuberger [BHN76] again studied the bosonization of $su(n)$ Thirring models, this time with periodic boundary conditions as $L \rightarrow \infty$. Translating the fermionic zero modes of their solution at fixed L into vertex-operators on the string, one now understands that integer-moded complex versions (CR) of world-sheet R fermions and $su(n)_1$ currents can also be obtained from the *same* vertex operators by choosing degenerate ground states with $\pm 1/2$ for each of the n momenta.

In later work (see for example Frenkel, Lepowsky and Meurman [FLM88] and Lepowsky [Lep07]), one learns that the true vacuum of the full CFT is the BH vacuum $|0\rangle = |\{p = J(0) = 0\}\rangle$ and products of BH currents on the BH vacuum live on the root-lattice of $su(n)$. The BH theory is itself reducible, containing many copies of the integrable representations of $su(n)$ – each copy living at different Δ 's on the weight lattice of $su(n)$.

1.8 The Confluence with Mathematics

So ended our three contributions [BH71], [Hal71c], [Hal75] to current-algebraic internal symmetry on the string. Loosely speaking, these papers also marked the end of the first string era, as an increasing number of physicists – including ourselves – turned their interest to gauge theory for a decade.

We were therefore fundamentally surprised to see in 1985 that the seminal paper on the heterotic string (Gross, Harvey, Martinec and Rohm [GHMR85]) used the 1980 vertex-operator (root-lattice) construction of $E(8)_1$ by Frenkel and Kac [FK80]. Following this, we contacted James Lepowsky and Igor Frenkel, who patiently exchanged information with us about these parallel developments in mathematics and physics, including explanations of the first vertex-operator construction (twisted $su(2)_1$) in mathematics (Lepowsky and Wilson [LW78]), and the earlier Kac-Moody algebras [Kac67], [Moo67] – which include the affine and hyperbolic algebras as special cases.

In mathematics, our 1971 dual quark-model construction of $su(3)_1$ is known as the *first concrete representation of affine Lie algebra*, including the first explicit central term. In this connection we quote first from page 6304 of a 1980 paper by Frenkel [Fre80]: “Ten years ago two physicists, Bardakci and Halpern, in Ref. [BH71] constructed a representation of the subalgebra $\tilde{gl}(l)$ of $\tilde{o}(2l)$ in the space $V((2\mathbb{Z} + 1)^l)$ (see formulas 3.1-3.11 of Ref. [BH71]). At that time the theory of affine Lie algebras began to take

its first steps.” A less technical form of this statement is found on page 36 of the Introduction to the review article by Frenkel, Lepowsky and Meurman [FLM88]: “In their proposal of current-algebraic internal symmetry for strings, Bardakci and Halpern [BH71], independently of mathematicians, discovered affine Lie algebras in 1971, including the first case of a concrete representation — a fermionic realization of $sl(\hat{3})$; irreducibility issues arose later in independent discoveries by mathematicians.”

The affine-Sugawara constructions ([BH71], [Hal71c], [KZ84]) in physics were also found in math [Seg81], where they are known as the Segal operators.

The early vertex-operator constructions in physics are also well-known in mathematics. In this connection, we find again on the same page of Ref. [FLM88]: “The untwisted vertex-operator construction of $sl(\hat{n})$ -representations from compactified spatial dimensions was implicit in [Hal75], [BHN76].” A supplementary statement is found on page 365 of Ref. [Lep07]: “It turned out that vertex operators and symmetry are closely related. Instances of this had already been discovered in physics, including the works [Hal75] and [BHN76].”

In conclusion, one might say that for us, ignorance was bliss – or, more optimistically – that physicists can create mathematics. In either case, it is clear that both fields have profited handsomely from this confluence.

1.9 The Larger Perspective

In our historical discussion above we have focused on the discovery of current-algebraic CFT on the string worldsheet, but to complete the modern perspective we should emphasize that this is just one of many mathematical methods which have been found and developed in string theory. Over the last 25 years, this domain has been the main source of new methods in theoretical physics that deeply changed the discipline and had many other applications in different areas, such as statistical mechanics and condensed matter physics. These developments involved the latest mathematical advances in algebra, topology and geometry. Some of these results (such as the affine algebras described here) were first discovered in string theory and then studied systematically by mathematicians, while others were imported from mathematics and made clear and practical by the string applications.

This communication and cross-fertilization has also had a large impact in mathematics, where the field theory methods are now well appreciated (see for example the lecture notes in Ref. [Princeton99]). In the 80’s and 90’s, Edward Witten has played a crucial role in this interdisciplinary endeavor.

In our context we note in particular the important set of string-theory mathematical methods given by general two-dimensional conformal field theory, which began with the analysis by Belavin, Polyakov and Zamolodchikov [BPZ84]. They used the mathematical results of Kac [Kac90] and Feigin and Fuchs [FF82] on representation theory of the Virasoro algebra, and realized their relevance for exact descriptions of two-dimensional critical phenomena (see for example the collection of important papers in [ISZ88]).

For help in the preparation of this article, we thank O. Ganor, I. Frenkel, Y. Frishman and J. Lepowsky.

References

- [AD68] Adler, S. and Dashen, R.F. (1968). *Current algebras* (W.A. Benjamin Inc., New York).
- [BHN76] Banks, T., Horn, D. and Neuberger, H. (1976). Bosonization of the SU(N) Thirring models, *Nucl. Phys.* **B108**, 119–129.
- [Bar71] Bardakci, K. (1971). New gauge identities in dual quark models, *Nucl. Phys.* **B33**, 464–474.
- [BFH68] Bardakci, K., Frishman, Y. and Halpern, M.B. (1968). Structure and extensions of a theory of currents, *Phys. Rev.* **170**, 1353–1359.
- [BH68] Bardakci, K. and Halpern, M.B. (1968). Canonical representation of Sugawara's theory, *Phys. Rev.* **172**, 1542–1550.
- [BH69] Bardakci, K. and Halpern, M.B. (1969). Possible Born term for the hadronic bootstrap, *Phys. Rev.* **183**, 1456–1462.
- [BH71] Bardakci, K. and Halpern, M.B. (1971). New dual quark models, *Phys. Rev.* **D3**, 2493–2506.
- [BPZ84] Belavin, A. A., Polyakov, A. M. and Zamolodchikov, A. B. (1984) Infinite conformal symmetry in two-dimensional quantum field theory *Nucl. Phys.* **B241**, 333.
- [CT69] Chan, H.M. and Tsou, S.T. (1969). Explicit construction of the N-point function in the generalized Veneziano model, *Phys. Lett.* **B28**, 485–488.
- [CT74] Chodos, A. and Thorn, C.B. (1974). Making the massless string massive, *Nucl. Phys.* **B72**, 509–522.
- [Col75] Coleman, S. (1975). Quantum sine-Gordon equation as the massive Thirring model, *Phys. Rev.* **D11**, 2088–2097.
- [CGJ69] Coleman, S., Gross, D. and Jackiw, R. (1969). Fermion avatars of the Sugawara model, *Phys. Rev.* **180**, 1359–1365.
- [DF73] Dashen, R.F. and Frishman, Y. (1973). Thirring model with U(n) symmetry - scale invariant only for fixed values of a coupling constant, *Phys. Lett.* **B46**, 439–442.
- [DF75] Dashen, R.F. and Frishman, Y. (1975). Four-fermion interactions and scale invariance, *Phys. Rev.* **D11**, 2781–2802.
- [Princeton99] Deligne, P., Etingof, P., Freed, D.S., Jeffrey, L., Kazhdan, L., Morgan, J., Morrison, D.R. and Witten, E., eds., (1999), *Quantum Fields and Strings: A Course For Mathematicians* 2 vols., (American Mathematical Society, Providence).
- [DFZ72] Dell'Antonio, G.F., Frishman, Y. and Zwanziger, D. (1974). Thirring model

- in terms of currents: Solutions and light-cone expansions, *Phys. Rev.* **D6**, 988–1007.
- [FF82] Feigin, B.L. and Fuchs, D.B. (1982). Verma modules over the Virasoro algebra, *Functional Analysis and its Applications* **17**, 241.
- [FH88] Freericks, J.K. and Halpern, M.B. (1988). Conformal deformation by the currents of affine g , *Ann. Phys.* **188**, 258–306.
- [Fre80] Frenkel, I.B. (1980). Spinor representations of affine Lie algebras, *Proc. Nat. Acad. Sci. USA* **77**, 6303–6306.
- [FK80] Frenkel, I.B. and Kac, V.G. (1980). Basic representations of affine Lie algebras and dual resonance models, *Inv. Math.* **62**, 23.
- [FLM88] Frenkel, I.B., Lepowsky, J. and Meurman, A. (1988). *Vertex operator algebras and the monster*, *Pure and Applied Math.*, Vol. 134 (Academic Press, Boston).
- [FV69] Fubini, S. and Veneziano, G. (1969). Level structure of dual resonance models, *Nuovo Cim.* **A64**, 811–840.
- [GW86] Gepner, D. and Witten, E. (1986) String Theory on Group Manifolds, *Nucl. Phys.* **B278**, 493.
- [GKO85] Goddard, P., Kent, A. and Olive, D. (1985). Virasoro algebras and coset models, *Phys. Lett.* **152B**, 88–92.
- [GS82] Green, M.B. and Schwarz, J.H. (1982). Supersymmetrical string theories, *Phys. Lett.* **B109**, 444–448.
- [GSW87] Green, M.B., Schwarz, J.H. and Witten, E. (1987). *Superstring theory* (Cambridge University Press, Cambridge).
- [GH69] Gross, D.J. and Halpern, M.B. (1969). Theory of currents and the non-strong interactions *Phys. Rev.* **179**, 1436–1444.
- [GHMR85] Gross, D.J., Harvey, J.A., Martinec, E. and Rohm, R. (1985). The heterotic string, *Phys. Rev. Lett.* **54**, 502–505.
- [Hal71a] Halpern, M.B. (1971). Persistence of the photon in conformal dual models, *Phys. Rev.* **D3**, 3068–3071.
- [Hal71b] Halpern, M.B. (1971). New dual models of pions with no tachyon, *Phys. Rev.* **D4**, 3082–3083.
- [Hal71c] Halpern, M.B. (1971). The two faces of a dual pion-quark model, *Phys. Rev.* **D4**, 2398–2401.
- [Hal75] Halpern, M.B. (1975). Quantum ‘solitons’ which are $SU(N)$ fermions, *Phys. Rev.* **D12**, 1684–1699.
- [Hal76] Halpern, M.B. (1976). Equivalent-Boson method and free currents in two-dimensional gauge theories, *Phys. Rev.* **D13**, 337–342.
- [HK89] Halpern, M.B. and Kiritsis, E. (1989). General Virasoro construction on affine g , *Mod. Phys. Lett.* **A4**, 1373–1380.
- [HKOC96] Halpern, M.B., Kiritsis, E., Obers, N.A. and Clubok, K. (1996). Irrational conformal field theory, *Phys. Rep.* **265**, 1–138. [hep-th/9501144](#).
- [HT71a] Halpern, M.B. and Thorn, C.B. (1971). Dual model of pions with no tachyon, *Phys. Lett.* **B35**, 441–442.
- [HT71b] Halpern, M.B. and Thorn, C.B. (1971). The two faces of a dual pion-quark model II. Fermions and other things, *Phys. Rev.* **D4**, 3084–3088.
- [ISZ88] Itzykson, C., Saleur, H. and Zuber, J.-B. (1988). *Conformal invariance and applications to statistical mechanics* (World Scientific, Singapore).
- [Kac67] Kac, V.G. (1967). Simple graded Lie algebras of finite growth, *Funct. Anal. Appl.* **1**, 328–329.
- [Kac90] Kac, V.G. (1990). *Infinite dimensional Lie algebras*, (Cambridge University

- Press, Cambridge).
- [Kle38] Klein, O. (1938). Approximate treatment of electrons in a crystal lattice, *J. Phys. Radium* **9**, 1–12.
 - [KZ84] Knizhnik, V.G. and Zamolodchikov, A.B. (1984). Current algebra and Wess-Zumino model in two dimensions, *Nucl. Phys.* **B247**, 83–103.
 - [KS75] Kogut, J. and Susskind, L. (1975). How quark confinement solves the $\eta_3 \rightarrow 3\pi$ problem, *Phys. Rev.* **D11**, 3594–3610.
 - [Lep07] Lepowsky, J. (2007). Some developments in vertex operator algebra theory, old and new, in: Lie algebras, vertex operator algebras and their applications, International Conference in honor of J. Lepowsky and R. Wilson, ed. by Y.-Z. Huang and K.C. Misra *Contemp. Math.*, Amer. Math. Soc., Vol. 442, p. 355.
 - [LW78] Lepowsky, J. and Wilson, R.L. (1978). Construction of the affine Lie algebra $A_1^{(1)}$, *Comm. Math. Phys.* **62**, 43–53.
 - [Man73a] Mandelstam, S. (1973). K-degeneracy in non-additive dual resonance models, *Phys. Rev.* **D7**, 3763–3776.
 - [Man73b] Mandelstam, S. (1973). Simple non-additive dual resonance model, *Phys. Rev.* **D7**, 3777–3784.
 - [Man74] Mandelstam, S. (1974). Dual resonance models, *Phys. Rep.* **C13**, 259–353.
 - [Man75] Mandelstam, S. (1975). Soliton operators for the quantized sine-Gordon equation, *Phys. Rev.* **D11**, 3026–3030.
 - [Moo67] Moody, R.V. (1967). Lie algebras associated with general Cartan matrices, *Bull. Am. Math. Soc.* **73**, 217–221.
 - [NS71] Neveu, A. and Schwarz, J.H. (1971). Factorizable dual model of pions, *Nucl. Phys.* **B31**, 86–112.
 - [PC69] Paton, J.E. and Chan, H.M. (1969). Generalized Veneziano model with isospin, *Nucl. Phys.* **B10**, 516–520.
 - [Ram71] Ramond, P. (1971). Dual theory of free fermions, *Phys. Rev.* **D3**, 2415–2418.
 - [Sch59] Schwinger, J. (1959). Field theory commutators, *Phys. Rev. Lett.* **3**, 296–297.
 - [Seg81] Segal, G. (1981). Unitary representations of some infinite dimensional groups, *Comm. Math. Phys.* **80**, 301–342.
 - [Som68] Sommerfield, C. (1968). Currents as dynamical variables, *Phys. Rev.* **176**, 2019–2025.
 - [Sug68] Sugawara, H. (1968). A field theory of currents, *Phys. Rev.* **170**, 1659–1662.
 - [Vir70] Virasoro, M. (1970). Subsidiary conditions and ghosts in dual-resonance models, *Phys. Rev.* **D1**, 2933–2936.
 - [Wit84] Witten, E. (1984) Nonabelian bosonization in two dimensions, *Commun. Math. Phys.* **92**, 455.